


Exposures negatively associated with uterine cancer risk

Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
카로티노이드				
루테인	3	3,315	1,227	0.5 (0.34-0.71)
베타-카로틴	3	3,315	1,227	0.6 (0.46-0.78)
라이코펜	3	3,463	987	0.92 (0.66-1.29)
비타민 A, C, D, E, K				
비타민 E	2	30,376	2,937	0.68 (0.37-1.24)
비타민 D	4	66,026	2,398	0.82 (0.59-1.14)
비타민 C	2	121,756	13,138	0.86 (0.36-2.07)
비타민 B군				
엽산	2	74,680	412	0.49 (0.32-0.73)
무기질 및 미량 원소				
칼슘	4	6,240	1,445	0.68 (0.55-0.84)
폴리페놀 및 플라보노이드				
녹차	3	3,331	1,564	0.7 (0.51-0.96)
차	4	4,413	2,105	0.76 (0.48-1.21)
지방산 및 지질				
오메가-3 지방산	5	457,863	4,240	0.86 (0.64-1.15)
어유	3	110,943	2,072	0.87 (0.48-1.61)
식물성 에스트로겐				
대두	3	95,991	933	0.73 (0.48-1.13)
기타				
지중해식 식단	3	90,098	3,100	0.63 (0.37-1.08)
카페인	4	150,854	1,222	0.91 (0.74-1.11)
알코올	19	3,080,855	192,818	0.95 (0.85-1.05)



Exposure group



비타민 B군



카로티노이드



지방산 및 지질



무기질 및 미량 원소



기타



식물성 에스트로겐



폴리페놀 및 플라보노이드



비타민 A, C, D, E, K