


Exposures negatively associated with uterine cancer risk

Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
类胡萝卜素				
叶黄素	3	3,315	1,227	0.5 (0.34–0.71)
β-胡萝卜素	3	3,315	1,227	0.6 (0.46–0.78)
番茄红素	3	3,463	987	0.92 (0.66–1.29)
维生素 A、C、D、E、K				
维生素 E	2	30,376	2,937	0.68 (0.37–1.24)
维生素 D	4	66,026	2,398	0.82 (0.59–1.14)
维生素 C	2	121,756	13,138	0.86 (0.36–2.07)
B族维生素				
叶酸	2	74,680	412	0.49 (0.32–0.73)
矿物质与微量元素				
钙	4	6,240	1,445	0.68 (0.55–0.84)
多酚与黄酮类				
绿茶	3	3,331	1,564	0.7 (0.51–0.96)
茶	4	4,413	2,105	0.76 (0.48–1.21)
脂肪酸与脂质				
Omega-3 脂肪酸	5	457,863	4,240	0.86 (0.64–1.15)
鱼油	3	110,943	2,072	0.87 (0.48–1.61)
植物雌激素				
大豆	3	95,991	933	0.73 (0.48–1.13)
其他				
地中海饮食	3	90,098	3,100	0.63 (0.37–1.08)
咖啡因	4	150,854	1,222	0.91 (0.74–1.11)
酒精	19	3,080,855	192,818	0.95 (0.85–1.05)



Exposure group

- B族维生素
- 类胡萝卜素
- 脂肪酸与脂质
- 矿物质与微量元素
- 其他
- 植物雌激素
- 多酚与黄酮类
- 维生素 A、C、D、E、K

RR < 1 = inversely associated with uterine cancer risk. | [*] pooled RR (size proportional to # of studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval