


Exposures positively associated with breast cancer risk meta-analysis of dietary-intake studies				
Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
維生素 A、C、D、E、K				
維生素 K	2	5,048	2,592	1.01 (0.68–1.5)
礦物質與微量元素				
硒	6	175,826	12,809	1.02 (0.87–1.2)
鐵	8	385,425	22,415	1.22 (1.04–1.43)
多酚與類黃酮				
茶	26	1,631,244	201,438	1.03 (0.96–1.12)
水果與蔬菜				
葡萄柚	2	164,504	5,404	1.1 (0.8–1.52)
脂肪酸與脂質				
共軛亞油酸	4	380,443	17,329	1.06 (0.92–1.22)
Omega-6 脂肪酸	2	2,476	1,201	1.74 (1.26–2.42)
其他				
雞蛋	8	29,311	5,421	1.19 (0.99–1.43)
酒精	185	16,045,202	1,276,703	1.24 (1.21–1.26)
紅肉	34	1,178,141	60,075	1.33 (1.24–1.42)



Exposure group

- 脂肪酸與脂質
- 水果與蔬菜
- 礦物質與微量元素
- 其他
- 多酚與類黃酮
- 維生素 A、C、D、E、K

RR > 1 = positively associated with breast cancer risk. | [*] pooled RR (size proportional to # of studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval