


Exposures inversely associated with ovarian cancer risk

Exposure	N studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	4	529,715	3,624	0.69 (0.48–0.98)
beta-Carotene	9	678,322	30,201	0.87 (0.78–0.98)
VITAMINS A, C, D, E, K				
Vitamin E	7	657,479	30,006	0.65 (0.48–0.88)
Vitamin C	3	583,089	2,528	0.72 (0.41–1.29)
Vitamin D	13	426,489	19,570	0.81 (0.73–0.91)
Vitamin A	6	1,588,466	9,351	0.97 (0.86–1.09)
B VITAMINS				
Folic Acid	9	712,037	10,452	0.87 (0.68–1.1)
MINERALS & TRACE ELEMENTS				
Selenium	3	1,947	768	0.23 (0.06–0.93)
Calcium	7	42,324	4,508	0.69 (0.53–0.9)
POLYPHENOLS & FLAVONOIDS				
Tea	6	238,872	3,158	0.77 (0.53–1.13)
FRUITS & VEGETABLES				
Legumes	2	63,479	506	0.6 (0.25–1.43)
FATTY ACIDS & LIPIDS				
Omega-6 FA	2	6,630	3,238	0.8 (0.69–0.91)
PHYTOESTROGENS				
Soy	2	1,906	754	0.41 (0.25–0.67)
OTHER				
Alcohol	18	2,268,288	41,778	0.96 (0.84–1.09)
Caffeine	12	380,116	39,052	0.97 (0.86–1.09)



Exposure group

-  B Vitamins
-  Carotenoids
-  Fatty Acids & Lipids
-  Fruits & Vegetables
-  Minerals & Trace Elements
-  Other
-  Phytoestrogens
-  Polyphenols & Flavonoids
-  Vitamins A, C, D, E, K

RR < 1 = inversely associated with ovarian cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval