


Exposures inversely associated with uterine cancer risk				
Exposure	N studies	N	Cases	Pooled RR (95% CI)
类胡萝卜素				
叶黄素	3	3,315	1,227	0.5 (0.34–0.71)
β-胡萝卜素	3	3,315	1,227	0.6 (0.46–0.78)
番茄红素	2	2,233	686	0.8 (0.54–1.19)
维生素 A、C、D、E、K				
维生素 C	2	121,756	13,138	0.86 (0.36–2.07)
维生素 D	5	66,026	3,042	0.88 (0.69–1.12)
B族维生素				
叶酸	2	74,680	412	0.49 (0.32–0.73)
矿物质与微量元素				
钙	4	6,240	1,445	0.68 (0.55–0.84)
多酚与黄酮类				
绿茶	3	3,331	1,564	0.7 (0.51–0.96)
茶	4	4,413	2,105	0.76 (0.48–1.21)
脂肪酸与脂质				
Omega-3 脂肪酸	5	457,863	4,240	0.86 (0.64–1.15)
鱼油	3	110,943	2,072	0.87 (0.48–1.61)
植物雌激素				
大豆	3	95,991	933	0.73 (0.48–1.13)
其他				
地中海饮食	3	90,098	3,100	0.63 (0.37–1.08)
酒精	18	2,958,970	189,040	0.88 (0.78–1)
咖啡因	4	150,854	1,222	0.91 (0.74–1.11)



Exposure group

B族维生素

类胡萝卜素

脂肪酸与脂质

矿物质与微量元素

其他

植物雌激素

多酚与黄酮类

维生素 A、C、D、E、K

RR < 1 = inversely associated with uterine cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval