


Exposures inversely associated with ovarian cancer risk
meta-analysis of dietary-intake studies

Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	4	529,715	3,624	0.69 (0.48–0.98)
beta-Carotene	8	611,872	4,692	0.74 (0.6–0.92)
VITAMINS A, C, D, E, K				
Vitamin D	2	4,611	1,993	0.55 (0.41–0.74)
Vitamin E	6	591,029	4,497	0.69 (0.51–0.93)
Vitamin C	3	583,089	2,528	0.72 (0.41–1.29)
Vitamin A	6	1,588,466	9,351	0.97 (0.86–1.09)
B VITAMINS				
Folic Acid	8	723,200	3,495	0.77 (0.58–1.03)
ANTIOXIDANTS				
Antioxidants	4	523,871	2,791	0.62 (0.39–0.99)
MINERALS & TRACE ELEMENTS				
Selenium	2	1,633	611	0.62 (0.41–0.92)
Calcium	3	36,513	1,667	0.69 (0.57–0.83)
POLYPHENOLS & FLAVONOIDS				
Tea	8	307,213	4,002	0.79 (0.58–1.07)
FRUITS & VEGETABLES				
Legumes	2	63,479	506	0.6 (0.25–1.43)
FATTY ACIDS & LIPIDS				
Omega-6 FA	2	6,630	3,238	0.8 (0.69–0.91)
PHYTOESTROGENS				
Soy	2	1,906	754	0.41 (0.25–0.67)
OTHER				
Alcohol	18	819,111	17,098	0.96 (0.84–1.11)
Caffeine	11	336,054	38,479	0.96 (0.8–1.14)



Exposure group

- Antioxidants
- B Vitamins
- Carotenoids
- Fatty Acids & Lipids
- Fruits & Vegetables
- Minerals & Trace Elements
- Other
- Phytoestrogens
- Polyphenols & Flavonoids
- Vitamins A, C, D, E, K

RR < 1 = inversely associated with ovarian cancer risk. | [*] pooled RR (size proportional to # of studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval