


Exposures negatively associated with uterine cancer risk meta-analysis of dietary-intake studies				
Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	3	3,315	1,227	0.47 (0.32–0.69)
beta-Carotene	3	3,315	1,227	0.57 (0.43–0.76)
Lycopene	2	2,233	686	0.78 (0.5–1.21)
B VITAMINS				
Folic Acid	2	74,680	412	0.47 (0.31–0.71)
MINERALS & TRACE ELEMENTS				
Calcium	4	6,240	1,445	0.66 (0.52–0.82)
POLYPHENOLS & FLAVONOIDS				
Green Tea	3	3,331	1,564	0.68 (0.48–0.95)
Tea	4	4,413	2,105	0.74 (0.45–1.21)
FATTY ACIDS & LIPIDS				
Omega-3 FA	5	457,863	4,240	0.85 (0.62–1.16)
Fish Oil	3	110,943	2,072	0.87 (0.46–1.64)
PHYTOESTROGENS				
Soy	3	95,991	933	0.73 (0.46–1.13)
OTHER				
Mediterranean Diet	3	90,098	3,100	0.61 (0.34–1.09)
Caffeine	4	150,854	1,222	0.9 (0.73–1.12)
Alcohol	15	411,289	8,229	0.96 (0.85–1.08)



Exposure group

-  B Vitamins
-  Carotenoids
-  Fatty Acids & Lipids
-  Minerals & Trace Elements
-  Other
-  Phytoestrogens
-  Polyphenols & Flavonoids