


Exposures negatively associated with uterine cancer risk
meta-analysis of dietary-intake studies

Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
類胡蘿蔔素				
葉黃素	3	3,315	1,227	0.47 (0.32–0.69)
β-胡蘿蔔素	3	3,315	1,227	0.57 (0.43–0.76)
茄紅素	2	2,233	686	0.78 (0.5–1.21)
B群維生素				
葉酸	2	74,680	412	0.47 (0.31–0.71)
礦物質與微量元素				
鈣	4	6,240	1,445	0.66 (0.52–0.82)
多酚與類黃酮				
綠茶	3	3,331	1,564	0.68 (0.48–0.95)
茶	4	4,413	2,105	0.74 (0.45–1.21)
脂肪酸與脂質				
Omega-3 脂肪酸	5	457,863	4,240	0.85 (0.62–1.16)
魚油	3	110,943	2,072	0.87 (0.46–1.64)
植物雌激素				
黃豆	3	95,991	933	0.73 (0.46–1.13)
其他				
地中海飲食	3	90,098	3,100	0.61 (0.34–1.09)
咖啡因	4	150,854	1,222	0.9 (0.73–1.12)
酒精	15	411,289	8,229	0.96 (0.85–1.08)



Exposure group

-  B群維生素
-  類胡蘿蔔素
-  脂肪酸與脂質
-  礦物質與微量元素
-  其他
-  植物雌激素
-  多酚與類黃酮