


Exposures negatively associated with uterine cancer risk
meta-analysis of dietary-intake studies

Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
類胡蘿蔔素				
葉黃素	3	3,315	1,227	0.5 (0.34–0.71)
β-胡蘿蔔素	3	3,315	1,227	0.6 (0.46–0.78)
茄紅素	2	2,233	686	0.8 (0.54–1.19)
B群維生素				
葉酸	2	74,680	412	0.49 (0.32–0.73)
礦物質與微量元素				
鈣	4	6,240	1,445	0.68 (0.55–0.84)
多酚與類黃酮				
綠茶	3	3,331	1,564	0.7 (0.51–0.96)
茶	4	4,413	2,105	0.76 (0.48–1.21)
脂肪酸與脂質				
Omega-3 脂肪酸	5	457,863	4,240	0.86 (0.64–1.15)
魚油	3	110,943	2,072	0.87 (0.48–1.61)
植物雌激素				
黃豆	3	95,991	933	0.73 (0.48–1.13)
其他				
地中海飲食	3	90,098	3,100	0.63 (0.37–1.08)
咖啡因	4	150,854	1,222	0.91 (0.74–1.11)
酒精	15	411,289	8,229	0.96 (0.85–1.08)



Exposure group

- B群維生素
- 類胡蘿蔔素
- 脂肪酸與脂質
- 礦物質與微量元素
- 其他
- 植物雌激素
- 多酚與類黃酮

RR < 1 = inversely associated with uterine cancer risk. | [*] pooled RR (size proportional to # of studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval