


Exposures inversely associated with uterine cancer risk
meta-analysis of dietary-intake studies

Exposure	N studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	3	3,315	1,227	0.5 (0.34–0.71)
beta-Carotene	3	3,315	1,227	0.6 (0.46–0.78)
Lycopene	2	2,233	686	0.8 (0.54–1.19)
B VITAMINS				
Folic Acid	2	74,680	412	0.49 (0.32–0.73)
MINERALS & TRACE ELEMENTS				
Calcium	4	6,240	1,445	0.68 (0.55–0.84)
POLYPHENOLS & FLAVONOIDS				
Green Tea	3	3,331	1,564	0.7 (0.51–0.96)
Tea	4	4,413	2,105	0.76 (0.48–1.21)
FATTY ACIDS & LIPIDS				
Omega-3 FA	5	457,863	4,240	0.86 (0.64–1.15)
Fish Oil	3	110,943	2,072	0.87 (0.48–1.61)
PHYTOESTROGENS				
Soy	3	95,991	933	0.73 (0.48–1.13)
OTHER				
Mediterranean Diet	3	90,098	3,100	0.63 (0.37–1.08)
Alcohol	15	411,289	8,229	0.91 (0.8–1.04)
Caffeine	4	150,854	1,222	0.91 (0.74–1.11)



Exposure group



RR < 1 = inversely associated with uterine cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval