


Exposures positively associated with breast cancer risk
meta-analysis of dietary-intake studies

Exposure	N studies	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin K	2	5,048	2,592	1.01 (0.68–1.5)
MINERALS & TRACE ELEMENTS				
Selenium	6	175,826	12,809	1.02 (0.87–1.2)
Iron	8	385,425	22,415	1.22 (1.04–1.43)
POLYPHENOLS & FLAVONOIDS				
Tea	26	1,631,244	201,438	1.03 (0.96–1.12)
FRUITS & VEGETABLES				
Grapefruit	2	164,504	5,404	1.1 (0.8–1.52)
FATTY ACIDS & LIPIDS				
Conj. LA	4	380,443	17,329	1.06 (0.92–1.22)
Omega-6 FA	2	2,476	1,201	1.74 (1.26–2.42)
OTHER				
Eggs	8	29,311	5,421	1.19 (0.99–1.43)
Alcohol	187	16,101,045	1,279,155	1.24 (1.22–1.27)
Red Meat	34	1,178,141	60,075	1.33 (1.24–1.42)



Exposure group

- Fatty Acids & Lipids
- Fruits & Vegetables
- Minerals & Trace Elements
- Other
- Polyphenols & Flavonoids
- Vitamins A, C, D, E, K

RR > 1 = positively associated with breast cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval