


Exposures positively associated with breast cancer risk
meta-analysis of dietary-intake studies

Exposure	N studies	N	Cases	Pooled RR (95% CI)
VITAMINEN A, C, D, E EN K				
Vitamine K	2	5,048	2,592	1.01 (0.68-1.5)
MINERALEN EN SPORENELEMENTEN				
Seleen	6	175,826	12,809	1.02 (0.87-1.2)
IJzer	8	385,425	22,415	1.22 (1.04-1.43)
POLYFENOLEN EN FLAVONOÏDEN				
Thee	26	1,631,244	201,438	1.03 (0.96-1.12)
FRUIT EN GROENTEN				
Grapefruit	2	164,504	5,404	1.1 (0.8-1.52)
VETZUREN EN LIPIDEN				
Geconjugeerd linolzuur		380,443	17,329	1.06 (0.92-1.22)
Omega 6-vetzuren	2	2,476	1,201	1.74 (1.26-2.42)
OVERIG				
Eieren	8	29,311	5,421	1.19 (0.99-1.43)
Alcohol	187	16,101,045	1,279,155	1.24 (1.22-1.27)
Rood vlees	34	1,178,141	60,075	1.33 (1.24-1.42)



Exposure group

-  Vetzuren en lipiden
-  Fruit en groenten
-  Mineralen en sporenelementen
-  Overig
-  Polyfenolen en flavonoïden
-  Vitaminen A, C, D, E en K

RR > 1 = positively associated with breast cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval