
# Exposures positively associated with aggressive prostate cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Calcium** | 5 | 98,670 | - | *1.62 (0.92-2.84)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Alpha-linolenic acid** | 3 | 383,989 | 3,504 | *1.85 (1.03-3.31)* |
| **Monounsaturated Fat (MUFA)** | 2 | 2,234 | - | *2.39 (1.37-4.15)* |
| ***DIET*** | | | | |
| **Mediterranean diet** | 2 | 106,146 | - | *1.02 (0.92-1.13)* |
| ***OTHER*** | | | | |
| **Multivitamin** | 2 | 343,481 | 2,040 | *1.27 (1.04-1.56)* |
| **Alcohol** | 4 | 166,503 | - | *1.28 (0.73-2.26)* |
| **Eggs** | 2 | 869,756 | 5,123 | *1.36 (0.87-2.10)* |
| **Dairy** | 3 | 101,926 | 744 | *1.41 (1.14-1.75)* |
| **Processed meat intake** | 4 | 204,990 | 2,302 | *1.41 (1.21-1.65)* |
| **Red meat** | 5 | 129,031 | 1,575 | *1.44 (1.13-1.84)* |




Exposure group
Minerals & Trace Elements
Fatty Acids & Lipids
Diet
Other

*RR > 1 = positively associated with aggressive prostate cancer risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis*

Include subexposures: No

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.