
Exposures positively associated with aggressive prostate cancer risk

Exposure	Studies	N	Cases	Pooled RR (95% CI)
MINERALS & TRACE ELEMENTS				
Calcium	6	98,931	-	*1.53 (0.98-2.38)*
FATTY ACIDS & LIPIDS				
Monounsaturated Fat (MUFA)	3	144,473	-	*1.72 (0.85-3.51)*
Alpha-linolenic acid	3	383,989	3,504	*1.85 (1.03-3.31)*
Omega-3 fatty acids	3	383,989	3,504	*1.85 (1.03-3.31)*
Polyunsaturated Fat (PUFA)	4	385,026	-	*1.97 (1.16-3.35)*
METABOLITES & AMINO ACIDS				
Betaine	3	55,116	963	*1.03 (0.60-1.74)*
DIET				
Mediterranean diet	2	106,146	-	*1.02 (0.92-1.13)*
OTHER				
Multivitamin	2	343,481	2,040	*1.27 (1.04-1.56)*
Alcohol	4	166,503	-	*1.28 (0.73-2.26)*
Eggs	2	869,756	5,123	*1.36 (0.87-2.10)*
Dairy	3	101,926	744	*1.41 (1.14-1.75)*
Processed meat intake	4	204,990	2,302	*1.41 (1.21-1.65)*
Red meat	5	129,031	1,575	*1.44 (1.13-1.84)*



RR > 1 = positively associated with aggressive prostate cancer risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis

Include subexposures: Yes

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.