
# Exposures positively associated with kidney and renal pelvis cancer risk

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Copper** | 2 | 95,958 | 353 | *2.44 (0.43-13.84)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Dietary Protein Intake** | 2 | 776,349 | 2,168 | *1.30 (0.76-2.24)* |
| ***OTHER*** | | | | |
| **Sugar-Sweetened Beverage Intake** | 3 | 563,999 | 1,503 | *1.11 (0.96-1.28)* |
| **Ultra-Processed Food Intake** | 2 | 582,035 | 3,740 | *1.22 (1.03-1.45)* |
| **Processed Meat Intake** | 3 | 991,425 | 2,649 | *1.23 (1.05-1.44)* |
| **Red Meat** | 4 | 986,845 | 3,043 | *1.75 (1.22-2.53)* |
| **Dairy** | 3 | 3,506 | 1,009 | *2.76 (1.23-6.20)* |




Exposure group

Minerals & Trace Elements  Metabolites & Amino Acids  Other

*RR > 1 = positively associated with kidney and renal pelvis cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Include subexposures: No