
Exposures negatively associated with renal cell carcinoma risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Beta-Carotene	5	881,922	4,765	*0.75 (0.64-0.90)*
Lutein	4	879,626	3,623	*0.80 (0.69-0.91)*
Lycopene	3	873,444	2,860	*0.84 (0.59-1.22)*
VITAMINS A, C, D, E, K				
Vitamin A	2	4,597	1,909	*0.65 (0.43-0.99)*
Vitamin E	4	127,855	2,404	*0.67 (0.45-0.99)*
Vitamin C	5	128,222	2,559	*0.76 (0.61-0.94)*
Vitamin D	5	79,250	2,416	*0.78 (0.68-0.90)*
B VITAMINS				
Vitamin B12	2	1,959	730	*0.63 (0.47-0.85)*
Folic Acid	6	132,098	3,299	*0.65 (0.48-0.89)*
Vitamin B1	2	4,597	1,909	*0.71 (0.55-0.92)*
Vitamin B6	3	3,861	1,547	*0.74 (0.40-1.40)*
Vitamin B2	3	3,861	1,547	*0.84 (0.68-1.04)*
MINERALS & TRACE ELEMENTS				
Selenium	2	3,471	1,543	*0.26 (0.08-0.92)*
Calcium	2	2,274	931	*0.62 (0.16-2.40)*
POLYPHENOLS & FLAVONOIDS				
Tea	4	804,643	2,861	*0.72 (0.52-1.01)*
OTHER				
Alcohol	16	3,497,583	8,299	*0.80 (0.74-0.86)*
Total Dietary Fiber Intake	2	4,410	1,621	*0.85 (0.64-1.12)*



RR < 1 = negatively associated with renal cell carcinoma risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No