
Exposures positively associated with kidney and renal pelvis cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin D	3	327,723	1,577	*1.15 (0.58-2.30)*
MINERALS & TRACE ELEMENTS				
Copper	2	95,958	353	*2.44 (0.43-13.84)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	776,349	2,168	*1.30 (0.76-2.24)*
OTHER				
Sugar-Sweetened Beverage Intake	3	563,999	1,503	*1.11 (0.96-1.28)*
Ultra-Processed Food Intake	2	582,035	3,740	*1.22 (1.03-1.45)*
Processed Meat Intake	3	991,425	2,649	*1.23 (1.05-1.44)*
Red Meat	4	986,845	3,043	*1.75 (1.22-2.53)*
Dairy	3	3,506	1,009	*2.76 (1.23-6.20)*



Exposure group

Vitamins A, C, D, E, K | Minerals & Trace Elements | Metabolites & Amino Acids | Other

RR > 1 = positively associated with kidney and renal pelvis cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No