
Exposures negatively associated with bladder cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin C	2	4,359	831	*0.68 (0.43-1.07)*
Vitamin E	3	36,672	978	*0.80 (0.60-1.05)*
B VITAMINS				
Vitamin B1	2	533,488	3,237	*0.91 (0.52-1.60)*
FRUITS & VEGETABLES				
Legumes	2	2,541	1,268	*0.64 (0.43-0.95)*
Avocado	2	112,328	110	*0.78 (0.60-1.01)*
FERMENTED FOODS & PROBIOTICS				
Fermented Foods	2	82,627	665	*0.62 (0.48-0.80)*
METABOLITES & AMINO ACIDS				
Plant Protein Intake	2	903,751	2,856	*0.82 (0.72-0.93)*
DIET				
DASH Diet	3	187,282	380	*0.61 (0.30-1.23)*
Mediterranean Diet	6	1,536,289	8,379	*0.86 (0.77-0.97)*
OTHER				
Whole-Grain Intake	3	576,212	3,892	*0.72 (0.52-1.01)*
Dairy	6	186,061	2,176	*0.78 (0.53-1.13)*
Total Dietary Fiber Intake	2	676,447	3,990	*0.85 (0.76-0.95)*
Alcohol	8	950,368	6,562	*0.94 (0.79-1.12)*



RR < 1 = negatively associated with bladder cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No