
# Exposures negatively associated with kidney and renal pelvis cancer risk

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Beta-Carotene** | 5 | 881,922 | 4,765 | *0.76 (0.64-0.90)* |
| **Lutein** | 4 | 879,626 | 3,623 | *0.80 (0.69-0.91)* |
| **Lycopene** | 3 | 873,444 | 2,860 | *0.84 (0.59-1.22)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin A** | 2 | 4,597 | 1,909 | *0.66 (0.44-0.99)* |
| **Vitamin E** | 4 | 127,855 | 2,404 | *0.67 (0.45-0.99)* |
| **Vitamin C** | 5 | 128,222 | 2,559 | *0.76 (0.61-0.94)* |
| **Vitamin D** | 8 | 510,406 | 3,672 | *0.85 (0.61-1.19)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B12** | 2 | 1,959 | 730 | *0.63 (0.47-0.85)* |
| **Folic Acid** | 6 | 132,098 | 3,299 | *0.65 (0.48-0.89)* |
| **Vitamin B1** | 2 | 4,597 | 1,909 | *0.71 (0.55-0.92)* |
| **Vitamin B6** | 3 | 3,861 | 1,547 | *0.75 (0.40-1.40)* |
| **Vitamin B2** | 3 | 3,861 | 1,547 | *0.84 (0.68-1.04)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Selenium** | 2 | 3,471 | 1,543 | *0.26 (0.08-0.92)* |
| **Calcium** | 2 | 2,274 | 931 | *0.62 (0.16-2.39)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Green Tea** | 2 | 103,012 | 536 | *0.51 (0.24-1.11)* |
| **Tea** | 5 | 907,106 | 3,147 | *0.73 (0.56-0.96)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Choline** | 2 | 616,865 | 2,286 | *0.94 (0.84-1.06)* |
| ***OTHER*** | | | | |
| **Alcohol** | 24 | 4,606,074 | 15,007 | *0.84 (0.76-0.93)* |
| **Total Dietary Fiber Intake** | 3 | 198,798 | 2,632 | *0.87 (0.78-0.97)* |




*RR < 1 = negatively associated with kidney and renal pelvis cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Include subexposures: No