
Exposures positively associated with bladder cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
MINERALS & TRACE ELEMENTS				
Phosphorus	2	1,136	520	*1.19 (0.47-3.01)*
POLYPHENOLS & FLAVONOIDS				
Tea	5	39,321	4,484	*1.05 (0.72-1.53)*
PHYTOESTROGENS				
Soy	3	49,061	222	*1.12 (0.31-4.00)*
METABOLITES & AMINO ACIDS				
Animal Protein Intake	2	903,751	2,856	*1.14 (1.03-1.26)*
OTHER				
Processed Meat Intake	4	335,240	3,001	*1.09 (0.96-1.24)*
Red Meat	8	1,417,470	6,936	*1.27 (1.08-1.48)*



Exposure group

Minerals & Trace Elements | Polyphenols & Flavonoids | Phytoestrogens | Metabolites & Amino Acids | Other

RR > 1 = positively associated with bladder cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: Yes