
Exposures positively associated with bladder cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lycopene	2	222,535	1,249	*1.05 (0.87-1.28)*
B VITAMINS				
Vitamin B1	3	575,001	3,737	*1.03 (0.88-1.20)*
MINERALS & TRACE ELEMENTS				
Phosphorus	2	1,147	522	*1.19 (0.47-3.01)*
PHYTOESTROGENS				
Soy	3	49,061	222	*1.12 (0.31-4.00)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	585,408	2,402	*1.12 (1.01-1.23)*
Animal Protein Intake	2	585,408	2,402	*1.15 (1.04-1.28)*
OTHER				
Processed Meat Intake	7	938,913	6,956	*1.09 (0.98-1.21)*
Red Meat	9	1,417,360	6,876	*1.32 (1.11-1.57)*
Eggs	2	1,995	355	*3.86 (2.65-5.62)*



RR > 1 = positively associated with bladder cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No