
Exposures positively associated with kidney and renal pelvis cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin A	2	4,597	1,909	*1.23 (0.53-2.84)*
B VITAMINS				
Vitamin B1	2	4,597	1,909	*1.07 (0.51-2.23)*
MINERALS & TRACE ELEMENTS				
Copper	2	95,958	353	*2.44 (0.43-13.84)*
FATTY ACIDS & LIPIDS				
Monounsaturated Fat MUFA	2	777,095	1,801	*1.39 (0.82-2.35)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	776,349	2,168	*1.30 (0.76-2.24)*
OTHER				
Sugar Sweetened Beverage Intake	3	563,999	1,503	*1.11 (0.96-1.28)*
Processed Meat Intake	6	1,766,660	5,212	*1.15 (0.94-1.41)*
Red Meat	6	1,768,446	5,800	*1.43 (1.13-1.82)*
Dairy	4	19,420	1,051	*2.22 (1.24-3.97)*



RR > 1 = positively associated with kidney and renal pelvis cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: Yes