
Exposures negatively associated with bladder cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	2	1,969	940	*0.52 (0.32-0.82)*
Beta Carotene	2	57,269	1,178	*0.70 (0.45-1.08)*
Lycopene	2	101,940	858	*0.95 (0.90-0.99)*
VITAMINS A, C, D, E, K				
Vitamin D	8	15,511	3,156	*0.60 (0.46-0.79)*
Vitamin C	2	4,359	831	*0.68 (0.44-1.07)*
Vitamin E	2	1,785	754	*0.70 (0.50-0.98)*
B VITAMINS				
Folic Acid	2	2,508	1,233	*0.65 (0.36-1.17)*
Vitamin B1	3	403,230	1,548	*0.97 (0.87-1.08)*
MINERALS & TRACE ELEMENTS				
Selenium	7	161,335	3,086	*0.76 (0.56-1.03)*
FRUITS & VEGETABLES				
Legumes	2	2,541	1,268	*0.64 (0.43-0.95)*
Avocado	2	112,328	1,165	*0.78 (0.60-1.01)*
FERMENTED FOODS & PROBIOTICS				
Yogurt	2	101,981	906	*0.65 (0.25-1.65)*
Fermented Foods	3	203,114	2,164	*0.66 (0.55-0.80)*
FATTY ACIDS & LIPIDS				
Omega-3 Fatty Acids	2	1,061	572	*0.31 (0.20-0.48)*
OTHER				
Whole Grain Intake	2	1,486	678	*0.55 (0.23-1.33)*
Dairy	8	306,953	3,788	*0.74 (0.54-1.01)*
Total Dietary Fiber Intake	2	676,447	3,990	*0.85 (0.77-0.95)*
Alcohol	6	490,657	5,696	*0.91 (0.77-1.08)*



RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata

Include subexposures: No