
# Exposures negatively associated with bladder cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin C** | 2 | 4,359 | 831 | *0.68 (0.44-1.07)* |
| **Vitamin E** | 2 | 1,785 | 754 | *0.70 (0.50-0.98)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B1** | 3 | 403,230 | 1,548 | *0.97 (0.87-1.08)* |
| ***FRUITS & VEGETABLES*** | | | | |
| **Legumes** | 2 | 2,541 | 1,268 | *0.64 (0.43-0.95)* |
| **Avocado** | 2 | 112,328 | 1,165 | *0.78 (0.60-1.01)* |
| ***FERMENTED FOODS & PROBIOTICS*** | | | | |
| **Yogurt** | 2 | 101,981 | 906 | *0.65 (0.25-1.65)* |
| **Fermented Foods** | 3 | 203,114 | 2,164 | *0.66 (0.55-0.80)* |
| ***OTHER*** | | | | |
| **Whole Grain Intake** | 2 | 1,486 | 678 | *0.55 (0.23-1.33)* |
| **Dairy** | 8 | 306,953 | 3,788 | *0.74 (0.54-1.01)* |
| **Total Dietary Fiber Intake** | 2 | 676,447 | 3,990 | *0.85 (0.77-0.95)* |
| **Alcohol** | 6 | 490,657 | 5,696 | *0.91 (0.77-1.08)* |




*RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata*

Include subexposures: No