
Exposures negatively associated with renal cell carcinoma risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Zeaxanthin	2	781,129	2,616	*0.79 (0.67-0.93)*
Lutein	4	879,626	3,623	*0.80 (0.70-0.91)*
Beta Carotene	5	881,922	4,765	*0.98 (0.70-1.36)*
VITAMINS A, C, D, E, K				
Vitamin E	4	127,855	2,404	*0.67 (0.46-0.99)*
Vitamin C	5	128,222	2,559	*0.76 (0.62-0.94)*
B VITAMINS				
Vitamin B1	2	4,597	1,909	*0.72 (0.55-0.92)*
POLYPHENOLS & FLAVONOIDS				
Tea	3	777,802	2,495	*0.65 (0.43-0.98)*
OTHER				
Alcohol	15	2,782,705	21,572	*0.71 (0.63-0.80)*
Total Dietary Fiber Intake	2	4,410	1,621	*0.85 (0.65-1.12)*



RR < 1 = negatively associated with renal cell carcinoma risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No