Exposures positively associated with renal cell carcinoma risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lycopene	3	873,444	2,860	*1.02 (0.73-1.41)*
VITAMINS A, C, D, E, K				
Vitamin A	2	4,597	1,909	*1.22 (0.53-2.80)*
B VITAMINS				
Folic Acid	3	129,691	2,345	*1.04 (0.70-1.54)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	776,349	2,168	*1.30 (0.76-2.22)*
OTHER				
Processed Meat Intake	3	497,805	1,843	*1.37 (0.75-2.52)*
Red Meat	4	986,845	3,043	*1.76 (1.22-2.54)*
Dairy	3	3,506	1,009	*2.79 (1.24-6.32)*



Exposure group

Carotenoids | Vitamins A, C, D, E, K | B Vitamins | Metabolites & Amino Acids | Other

RR > 1 = positively associated with renal cell carcinoma risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: Yes