
Exposures negatively associated with bladder cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin C	2	4,359	831	*0.68 (0.44-1.07)*
Vitamin E	2	1,785	754	*0.70 (0.50-0.98)*
B VITAMINS				
Vitamin B1	3	403,230	1,548	*0.97 (0.87-1.08)*
FRUITS & VEGETABLES				
Legumes	2	2,541	1,268	*0.64 (0.43-0.95)*
Avocado	2	112,328	1,165	*0.78 (0.60-1.01)*
FERMENTED FOODS & PROBIOTICS				
Yogurt	2	101,981	906	*0.65 (0.25-1.65)*
Fermented Foods	3	203,114	2,164	*0.66 (0.55-0.80)*
OTHER				
Whole Grain Intake	2	1,486	678	*0.55 (0.23-1.33)*
Dairy	8	306,953	3,788	*0.74 (0.54-1.01)*
Total Dietary Fiber Intake	2	676,447	3,990	*0.85 (0.77-0.95)*
Alcohol	6	490,657	5,696	*0.91 (0.77-1.08)*



RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata

Include subexposures: Yes