
Exposures positively associated with kidney and renal pelvis cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin A	2	4,597	1,909	*1.23 (0.53-2.84)*
B VITAMINS				
Vitamin B1	2	4,597	1,909	*1.07 (0.51-2.23)*
MINERALS & TRACE ELEMENTS				
Copper	2	95,958	353	*2.44 (0.43-13.84)*
FATTY ACIDS & LIPIDS				
Monounsaturated Fat MUFA	2	777,095	1,801	*1.39 (0.82-2.35)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	776,349	2,168	*1.30 (0.76-2.24)*
OTHER				
Sugar Sweetened Beverage Intake	4	599,592	1,649	*1.13 (0.98-1.30)*
Processed Meat Intake	6	1,766,660	5,212	*1.15 (0.94-1.41)*
Ultra Processed Food Intake	3	1,127,884	4,245	*1.18 (1.08-1.30)*
Non Nutritive Sweetener Intake	2	115,981	473	*1.28 (1.00-1.63)*
Red Meat	6	1,768,446	5,800	*1.43 (1.13-1.82)*
Dairy	6	512,230	2,409	*1.57 (1.10-2.25)*



RR > 1 = positively associated with kidney and renal pelvis cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata

Include subexposures: Yes