
Exposures negatively associated with bladder cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Beta Carotene	5	366,541	1,449	*0.88 (0.71-1.10)*
VITAMINS A, C, D, E, K				
Vitamin C	3	8,606	902	*0.75 (0.56-1.01)*
Vitamin A	2	121,519	737	*0.77 (0.39-1.51)*
Vitamin E	4	157,524	1,453	*0.86 (0.69-1.07)*
B VITAMINS				
Folic Acid	4	307,597	1,465	*0.73 (0.51-1.05)*
MINERALS & TRACE ELEMENTS				
Phosphorus	2	1,147	522	*0.97 (0.27-3.50)*
POLYPHENOLS & FLAVONOIDS				
Tea	9	580,076	9,039	*0.96 (0.79-1.17)*
FRUITS & VEGETABLES				
Legumes	2	2,541	1,268	*0.64 (0.43-0.95)*
Avocado	2	112,328	1,165	*0.78 (0.60-1.01)*
FERMENTED FOODS & PROBIOTICS				
Yogurt	2	101,981	906	*0.64 (0.25-1.66)*
Fermented Foods	5	203,999	2,474	*0.75 (0.50-1.12)*
FATTY ACIDS & LIPIDS				
Monounsaturated Fat MUFA	2	648,914	3,592	*0.95 (0.44-2.05)*
METABOLITES & AMINO ACIDS				
Plant Protein Intake	2	585,408	2,402	*0.84 (0.70-1.00)*
OTHER				
Whole Grain Intake	3	219,042	1,666	*0.72 (0.52-1.01)*
Dairy	10	799,763	5,469	*0.83 (0.65-1.05)*
Total Dietary Fiber Intake	2	676,447	3,990	*0.85 (0.76-0.95)*
Alcohol	10	881,889	8,536	*0.86 (0.74-1.00)*



RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No