
Exposures positively associated with aggressive prostate cancer risk
Meta-analysis of dietary-intake studies

Exposure	Studies	N	Cases	Pooled RR (95% CI)
MINERALS & TRACE ELEMENTS				
Calcium	5	98,670	-	*1.62 (0.92-2.84)*
FATTY ACIDS & LIPIDS				
Alpha-linolenic acid	3	383,989	3,504	*1.85 (1.03-3.31)*
Monounsaturated Fat (MUFA)	2	2,234	-	*2.39 (1.37-4.15)*
Polyunsaturated Fat (PUFA)	6	721,149	-	*2.70 (1.10-6.80)*
Omega-3 fatty acids	5	720,112	4,355	*3.43 (1.67-7.04)*
DIET				
Mediterranean diet	2	106,146	-	*1.02 (0.92-1.13)*
OTHER				
Multivitamin	2	343,481	2,040	*1.27 (1.04-1.56)*
Alcohol	4	166,503	-	*1.28 (0.73-2.26)*
Dairy	2	80,266	439	*1.39 (1.08-1.78)*
Processed meat intake	3	177,383	2,103	*1.41 (1.20-1.65)*
Red meat	4	101,424	1,376	*1.57 (1.20-2.05)*



Exposure group
Minerals & Trace Elements
Fatty Acids & Lipids
Diet
Other

RR > 1 = positively associated with aggressive prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.