
# Exposures negatively associated with prostate cancer risk

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Zeaxanthin** | 4 | 2,106 | 808 | *0.66 (0.40-1.08)* |
| **Lycopene** | 20 | 321,509 | 28,185 | *0.85 (0.78-0.93)* |
| **Beta-carotene** | 15 | 122,141 | - | *0.92 (0.76-1.11)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin K** | 4 | 34,701 | 1,054 | *0.73 (0.58-0.92)* |
| **Vitamin C** | 11 | 68,944 | 7,699 | *0.87 (0.75-1.03)* |
| **Vitamin E** | 25 | 404,649 | 36,660 | *0.94 (0.87-1.02)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B6** | 3 | 70,879 | 7,656 | *0.89 (0.83-0.96)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Selenium** | 28 | 382,643 | 29,947 | *0.77 (0.67-0.89)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Green tea** | 7 | 99,090 | 1,577 | *0.62 (0.43-0.89)* |
| **Tea** | 10 | 314,398 | 15,389 | *0.75 (0.64-0.89)* |
| ***FRUITS & VEGETABLES*** | | | | |
| **Garlic** | 3 | 36,604 | 2,168 | *0.69 (0.42-1.13)* |
| **Tomato** | 11 | 1,057,923 | 67,108 | *0.74 (0.64-0.85)* |
| **Legumes** | 8 | 1,155,076 | 79,072 | *0.89 (0.81-0.97)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Linoleic acid** | 7 | 8,735 | 3,057 | *0.71 (0.50-0.99)* |
| **Oleic acid** | 2 | 58,714 | 851 | *0.80 (0.24-2.64)* |
| **Fish oil** | 2 | 37,289 | 1,735 | *0.89 (0.68-1.17)* |
| **Docosahexaenoic acid** | 3 | 144,141 | 7,729 | *0.94 (0.74-1.20)* |
| **Omega-6 fatty acids** | 11 | 264,882 | 5,940 | *0.96 (0.79-1.17)* |
| **Eicosapentaenoic acid** | 5 | 444,156 | 13,552 | *0.99 (0.89-1.11)* |
| ***PHYTOESTROGENS*** | | | | |
| **Soy** | 8 | 85,059 | 5,399 | *0.60 (0.41-0.89)* |
| **Genistein** | 9 | 91,302 | 3,818 | *0.72 (0.54-0.95)* |
| ***HERBAL & BOTANICAL*** | | | | |
| **Ginkgo biloba** | 2 | 38,308 | 1,665 | *0.93 (0.68-1.27)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Dietary protein intake** | 2 | 109,557 | 3,064 | *0.99 (0.89-1.11)* |
| ***HORMONES & ENDOGENOUS*** | | | | |
| **Melatonin** | 3 | 4,618 | 1,414 | *0.75 (0.45-1.23)* |
| ***DIET*** | | | | |
| **Plant-based diet** | 2 | 211,318 | 24,516 | *0.91 (0.84-0.99)* |
| **DASH Diet** | 2 | 152,714 | - | *0.93 (0.82-1.06)* |
| ***OTHER*** | | | | |
| **Insoluble dietary fiber intake** | 3 | 101,084 | - | *0.62 (0.38-1.03)* |
| **Soluble dietary fiber intake** | 3 | 80,888 | 10,851 | *0.89 (0.83-0.96)* |
| **Total dietary fiber intake** | 7 | 972,077 | 64,899 | *0.92 (0.86-1.00)* |
| **Eggs** | 3 | 900,752 | 53,381 | *0.99 (0.96-1.02)* |




Exposure group

Carotenoids
Vitamins A, C, D, E, K
B Vitamins
Minerals & Trace Elements
Polyphenols & Flavonoids
Fruits & Vegetables
Fatty Acids & Lipids
Phytoestrogens
Herbal & Botanical
Metabolites & Amino Acids
Hormones & Endogenous
Diet
Other

*RR < 1 = negatively associated with prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis*