
# Exposures positively associated with aggressive prostate cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Calcium** | 5 | 98,670 | - | *1.62 (0.92-2.84)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Alpha-linolenic acid** | 3 | 383,989 | 3,504 | *1.85 (1.03-3.31)* |
| **Omega-3 fatty acids** | 3 | 383,989 | 3,504 | *1.85 (1.03-3.31)* |
| **Polyunsaturated Fat (PUFA)** | 4 | 385,026 | - | *1.97 (1.16-3.35)* |
| **Monounsaturated Fat (MUFA)** | 2 | 2,234 | - | *2.39 (1.37-4.15)* |
| ***DIET*** | | | | |
| **Mediterranean diet** | 2 | 106,146 | - | *1.02 (0.92-1.13)* |
| ***OTHER*** | | | | |
| **Multivitamin** | 2 | 343,481 | 2,040 | *1.27 (1.04-1.56)* |
| **Alcohol** | 4 | 166,503 | - | *1.28 (0.73-2.26)* |
| **Dairy** | 2 | 80,266 | 439 | *1.39 (1.08-1.78)* |
| **Processed meat intake** | 3 | 177,383 | 2,103 | *1.41 (1.20-1.65)* |
| **Red meat** | 4 | 101,424 | 1,376 | *1.57 (1.20-2.05)* |




Exposure group

Minerals & Trace Elements  Fatty Acids & Lipids  Diet  Other

*RR > 1 = positively associated with aggressive prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis*

Include subexposures: Yes

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.