
Exposures negatively associated with kidney and renal pelvis cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Beta Carotene	5	881,922	4,765	*0.76 (0.64-0.90)*
Lutein	5	881,922	4,765	*0.77 (0.68-0.87)*
Zeaxanthin	2	781,129	2,616	*0.79 (0.67-0.93)*
Lycopene	3	873,444	2,860	*0.84 (0.59-1.22)*
VITAMINS A, C, D, E, K				
Vitamin E	4	127,855	2,404	*0.67 (0.45-0.99)*
Vitamin C	5	128,222	2,559	*0.76 (0.61-0.94)*
Vitamin D	12	584,968	5,299	*0.84 (0.65-1.08)*
B VITAMINS				
Vitamin B12	2	1,959	730	*0.63 (0.47-0.85)*
Folic Acid	6	84,212	3,088	*0.65 (0.47-0.89)*
Vitamin B6	3	3,861	1,547	*0.75 (0.40-1.40)*
Vitamin B2	3	3,861	1,547	*0.84 (0.68-1.04)*
MINERALS & TRACE ELEMENTS				
Selenium	2	3,471	1,543	*0.26 (0.08-0.92)*
Calcium	4	495,083	1,521	*0.85 (0.58-1.25)*
POLYPHENOLS & FLAVONOIDS				
Green Tea	2	103,012	536	*0.51 (0.24-1.11)*
Tea	5	907,106	3,147	*0.73 (0.56-0.95)*
METABOLITES & AMINO ACIDS				
Choline	2	568,979	2,075	*0.94 (0.83-1.06)*
OTHER				
Alcohol	28	14,155,769	29,965	*0.77 (0.70-0.85)*
Total Dietary Fiber Intake	3	198,798	2,632	*0.87 (0.78-0.97)*



RR < 1 = negatively associated with kidney and renal pelvis cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: No