
# Exposures negatively associated with renal cell carcinoma risk

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Lutein** | 5 | 881,922 | 4,765 | *0.77 (0.68-0.87)* |
| **Zeaxanthin** | 2 | 781,129 | 2,616 | *0.79 (0.67-0.93)* |
| **Beta Carotene** | 5 | 881,922 | 4,765 | *0.98 (0.70-1.36)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin E** | 4 | 127,855 | 2,404 | *0.67 (0.45-0.99)* |
| **Vitamin C** | 5 | 128,222 | 2,559 | *0.76 (0.61-0.94)* |
| **Vitamin D** | 9 | 153,812 | 2,840 | *0.82 (0.67-1.00)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B12** | 2 | 1,959 | 730 | *0.63 (0.47-0.85)* |
| **Vitamin B1** | 2 | 4,597 | 1,909 | *0.71 (0.55-0.92)* |
| **Vitamin B6** | 4 | 4,771 | 2,002 | *0.73 (0.52-1.04)* |
| **Folic Acid** | 6 | 84,212 | 3,088 | *0.76 (0.50-1.17)* |
| **Vitamin B2** | 3 | 3,861 | 1,547 | *0.84 (0.68-1.04)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Selenium** | 2 | 3,471 | 1,543 | *0.53 (0.04-7.15)* |
| **Calcium** | 2 | 2,273 | 154 | *0.62 (0.16-2.40)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Tea** | 4 | 804,643 | 2,861 | *0.71 (0.51-1.00)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Polyunsaturated Fat PUFA** | 3 | 861,316 | 2,507 | *0.94 (0.52-1.71)* |
| ***OTHER*** | | | | |
| **Alcohol** | 20 | 3,783,725 | 11,677 | *0.72 (0.65-0.79)* |
| **Total Dietary Fiber Intake** | 2 | 4,410 | 1,621 | *0.85 (0.64-1.12)* |




*RR < 1 = negatively associated with renal cell carcinoma risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Include subexposures: Yes