
Exposures positively associated with bladder cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
B VITAMINS				
Vitamin B1	3	575,001	3,737	*1.03 (0.88-1.20)*
MINERALS & TRACE ELEMENTS				
Calcium	3	493,396	1,881	*1.17 (0.83-1.64)*
METABOLITES & AMINO ACIDS				
Dietary Protein Intake	2	585,408	2,402	*1.12 (1.01-1.23)*
Animal Protein Intake	2	585,408	2,402	*1.15 (1.04-1.28)*
OTHER				
Ultra Processed Food Intake	2	647,537	1,906	*1.07 (0.91-1.26)*
Processed Meat Intake	8	939,835	6,986	*1.10 (0.99-1.22)*
Red Meat	9	1,417,360	6,876	*1.32 (1.11-1.57)*
Eggs	2	1,995	355	*3.86 (2.65-5.62)*



Exposure group

B Vitamins · Minerals & Trace Elements · Metabolites & Amino Acids · Other

RR > 1 = positively associated with bladder cancer risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Include subexposures: Yes