
# Exposures positively associated with kidney and renal pelvis cancer risk

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin A** | 2 | 4,597 | 1,909 | *1.23 (0.53-2.84)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B1** | 2 | 4,597 | 1,909 | *1.07 (0.51-2.23)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Copper** | 2 | 95,958 | 353 | *2.44 (0.43-13.84)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Dietary Protein Intake** | 2 | 776,349 | 2,168 | *1.30 (0.76-2.24)* |
| ***OTHER*** | | | | |
| **Sugar Sweetened Beverage Intake** | 4 | 599,592 | 1,649 | *1.13 (0.98-1.30)* |
| **Processed Meat Intake** | 6 | 1,766,660 | 5,212 | *1.15 (0.94-1.41)* |
| **Ultra Processed Food Intake** | 3 | 1,127,884 | 4,245 | *1.18 (1.08-1.30)* |
| **Non Nutritive Sweetener Intake** | 2 | 115,981 | 473 | *1.28 (1.00-1.63)* |
| **Red Meat** | 6 | 1,768,446 | 5,800 | *1.43 (1.13-1.82)* |
| **Dairy** | 6 | 512,230 | 2,409 | *1.57 (1.10-2.25)* |




Exposure group

Vitamins A, C, D, E, K | B Vitamins | Minerals & Trace Elements | Metabolites & Amino Acids | Other

*RR > 1 = positively associated with kidney and renal pelvis cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata*

Include subexposures: No