
# Exposures negatively associated with kidney and renal pelvis cancer risk

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Beta Carotene** | 5 | 881,922 | 4,765 | *0.76 (0.64-0.90)* |
| **Lutein** | 5 | 881,922 | 4,765 | *0.77 (0.68-0.87)* |
| **Zeaxanthin** | 2 | 781,129 | 2,616 | *0.79 (0.67-0.93)* |
| **Lycopene** | 3 | 873,444 | 2,860 | *0.84 (0.59-1.22)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin E** | 4 | 127,855 | 2,404 | *0.67 (0.45-0.99)* |
| **Vitamin C** | 5 | 128,222 | 2,559 | *0.76 (0.61-0.94)* |
| **Vitamin D** | 12 | 584,968 | 5,299 | *0.84 (0.65-1.08)* |
| ***B VITAMINS*** | | | | |
| **Vitamin B12** | 2 | 1,959 | 730 | *0.63 (0.47-0.85)* |
| **Folic Acid** | 6 | 84,212 | 3,088 | *0.65 (0.47-0.89)* |
| **Vitamin B6** | 3 | 3,861 | 1,547 | *0.75 (0.40-1.40)* |
| **Vitamin B2** | 3 | 3,861 | 1,547 | *0.84 (0.68-1.04)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Selenium** | 2 | 3,471 | 1,543 | *0.26 (0.08-0.92)* |
| **Calcium** | 4 | 495,083 | 1,521 | *0.85 (0.58-1.25)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Green Tea** | 2 | 103,012 | 536 | *0.51 (0.24-1.11)* |
| **Tea** | 5 | 907,106 | 3,147 | *0.73 (0.56-0.95)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Choline** | 2 | 568,979 | 2,075 | *0.94 (0.83-1.06)* |
| ***OTHER*** | | | | |
| **Alcohol** | 29 | 14,233,029 | 30,214 | *0.77 (0.70-0.84)* |
| **Total Dietary Fiber Intake** | 3 | 198,798 | 2,632 | *0.87 (0.78-0.97)* |




*RR < 1 = negatively associated with kidney and renal pelvis cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Include subexposures: No