
Reviewed preprint recoveries included

Exposures negatively associated with prostate cancer\aggressive risk

Exposure	Studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Zeaxanthin	4	2,106	808	*0.66 (0.40-1.08)*
Lycopene	16	146,572	17,462	*0.84 (0.74-0.94)*
Beta-carotene	13	91,971	7,275	*0.91 (0.73-1.12)*
VITAMINS A, C, D, E, K				
Vitamin C	11	68,944	7,699	*0.87 (0.75-1.03)*
Vitamin E	17	216,480	25,755	*0.93 (0.85-1.03)*
B VITAMINS				
Vitamin B6	3	70,879	7,656	*0.89 (0.83-0.96)*
MINERALS & TRACE ELEMENTS				
Selenium	22	223,759	27,039	*0.71 (0.59-0.87)*
POLYPHENOLS & FLAVONOIDS				
Green tea	6	49,170	1,463	*0.64 (0.42-0.96)*
Tea	8	217,811	12,238	*0.72 (0.58-0.90)*
FRUITS & VEGETABLES				
Garlic	3	36,604	2,168	*0.69 (0.42-1.13)*
Tomato	9	962,664	63,815	*0.73 (0.62-0.87)*
Legumes	7	1,072,593	74,668	*0.88 (0.78-0.98)*
FATTY ACIDS & LIPIDS				
Linoleic acid	7	8,735	3,057	*0.71 (0.50-0.99)*
Oleic acid	2	58,714	851	*0.80 (0.24-2.64)*
Fish oil	2	37,289	1,735	*0.89 (0.68-1.17)*
Docosahexaenoic acid	3	144,141	7,729	*0.94 (0.74-1.20)*
Omega-6 fatty acids	2	254,243	1,931	*0.96 (0.79-1.17)*
Eicosapentaenoic acid	5	444,156	13,552	*0.99 (0.89-1.11)*
PHYTOESTROGENS				
Soy	8	85,059	5,399	*0.60 (0.41-0.89)*
Genistein	7	62,306	2,581	*0.62 (0.45-0.85)*
HERBAL & BOTANICAL				
Ginkgo biloba	2	38,308	1,665	*0.93 (0.68-1.27)*
METABOLITES & AMINO ACIDS				
Dietary protein intake	2	109,557	3,064	*0.99 (0.89-1.11)*
HORMONES & ENDOGENOUS				
Melatonin	3	4,618	1,414	*0.75 (0.45-1.23)*
DIET				
Plant-based diet	2	211,318	24,516	*0.91 (0.84-0.99)*
OTHER				
Insoluble dietary fiber intake	2	57,649	6,553	*0.68 (0.38-1.23)*
Soluble dietary fiber intake	2	26,552	4,437	*0.89 (0.81-0.97)*
Total dietary fiber intake	6	917,741	58,485	*0.92 (0.84-1.02)*
Eggs	3	900,752	53,381	*0.99 (0.96-1.02)*



Exposure group

Carotenoids
Vitamins A, C, D, E, K
B Vitamins
Minerals & Trace Elements
Polyphenols & Flavonoids
Fruits & Vegetables
Fatty Acids & Lipids
Phytoestrogens
Herbal & Botanical
Metabolites & Amino Acids
Hormones & Endogenous
Diet
Other

RR < 1 = negatively associated with prostate cancer\aggressive risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | arrowhead = interval continues beyond the axis

Latest publication per cohort

Include subexposures: No