
# Exposures negatively associated with aggressive prostate cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin E** | 2 | 83,022 | 355 | *0.76 (0.29-1.94)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Zinc** | 2 | 82,482 | - | *0.84 (0.17-4.29)* |
| ***FRUITS & VEGETABLES*** | | | | |
| **Tomato** | 3 | 124,620 | - | *0.70 (0.54-0.90)* |
| ***OTHER*** | | | | |
| **Total dietary fiber intake** | 2 | 885,584 | 5,076 | *0.86 (0.63-1.17)* |




Exposure group

Vitamins A, C, D, E, K | Minerals & Trace Elements | Fruits & Vegetables | Other

*RR < 1 = negatively associated with aggressive prostate cancer risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available)*

Include subexposures: Yes

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.