
Exposures negatively associated with kidney and renal pelvis cancer risk
Meta-analysis of dietary-intake studies

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Beta Carotene	5	881,922	4,765	*0.76 (0.64-0.90)*
Lutein	5	881,922	4,765	*0.77 (0.68-0.87)*
Zeaxanthin	2	781,129	2,616	*0.79 (0.67-0.93)*
Lycopene	3	873,444	2,860	*0.84 (0.59-1.22)*
VITAMINS A, C, D, E, K				
Vitamin E	4	127,855	2,404	*0.67 (0.45-0.99)*
Vitamin C	5	128,222	2,559	*0.76 (0.61-0.94)*
Vitamin D	2	4,112	776	*0.79 (0.66-0.95)*
B VITAMINS				
Folic Acid	2	4,597	1,909	*0.74 (0.52-1.05)*
POLYPHENOLS & FLAVONOIDS				
Green Tea	2	103,012	536	*0.51 (0.24-1.11)*
Tea	5	907,106	3,147	*0.73 (0.56-0.95)*
OTHER				
Alcohol	26	13,972,515	29,505	*0.75 (0.68-0.83)*
Total Dietary Fiber Intake	3	198,798	2,632	*0.87 (0.78-0.97)*



Exposure group
Carotenoids · Vitamins A, C, D, E, K · B Vitamins · Polyphenols & Flavonoids · Other

RR < 1 = negatively associated with kidney and renal pelvis cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Latest publication per cohort

Include subexposures: No