
Reviewed preprint recoveries included

Exposures negatively associated with prostate cancer\aggressive risk
Meta-analysis of dietary-intake studies

Exposure	Studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	2	1,623	677	*0.65 (0.49-0.86)*
Beta-carotene	9	93,912	4,718	*0.80 (0.63-1.01)*
Lycopene	10	137,731	13,620	*0.86 (0.73-1.02)*
VITAMINS A, C, D, E, K				
Vitamin C	10	68,446	7,637	*0.86 (0.73-1.01)*
Vitamin E	10	176,954	12,118	*0.97 (0.88-1.08)*
B VITAMINS				
Vitamin B6	3	70,879	7,656	*0.89 (0.83-0.96)*
Folic acid	4	65,550	6,813	*0.91 (0.71-1.16)*
POLYPHENOLS & FLAVONOIDS				
Green tea	6	49,170	1,463	*0.64 (0.42-0.96)*
Tea	8	217,811	12,238	*0.72 (0.58-0.90)*
FRUITS & VEGETABLES				
Garlic	3	36,604	2,168	*0.69 (0.42-1.13)*
Tomato	9	962,664	63,815	*0.73 (0.62-0.87)*
Legumes	7	1,072,593	74,668	*0.88 (0.78-0.98)*
FATTY ACIDS & LIPIDS				
Eicosapentaenoic acid	2	430,507	7,761	*0.95 (0.79-1.15)*
Omega-3 fatty acids	3	98,585	8,483	*0.96 (0.85-1.08)*
PHYTOESTROGENS				
Genistein	2	43,907	440	*0.54 (0.37-0.80)*
Soy	8	85,059	5,399	*0.60 (0.41-0.89)*
HERBAL & BOTANICAL				
Ginkgo biloba	2	38,308	1,665	*0.93 (0.68-1.27)*
METABOLITES & AMINO ACIDS				
Dietary protein intake	2	109,557	3,064	*0.99 (0.89-1.11)*
DIET				
Plant-based diet	2	211,318	24,516	*0.91 (0.84-0.99)*
OTHER				
Insoluble dietary fiber intake	2	57,649	6,553	*0.68 (0.38-1.23)*
Soluble dietary fiber intake	2	26,552	4,437	*0.89 (0.81-0.97)*
Total dietary fiber intake	6	917,741	58,485	*0.92 (0.84-1.02)*
Eggs	3	900,752	53,381	*0.99 (0.96-1.02)*



RR < 1 = negatively associated with prostate cancer\aggressive risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available)

Latest publication per cohort

Include subexposures: No