
Reviewed preprint recoveries included

Exposures negatively associated with prostate cancer risk
Meta-analysis of dietary-intake studies

Exposure	Studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	2	1,623	677	*0.65 (0.49-0.86)*
Beta-carotene	10	94,949	-	*0.79 (0.63-0.98)*
Lycopene	13	282,888	22,602	*0.88 (0.79-0.98)*
VITAMINS A, C, D, E, K				
Vitamin C	10	68,446	7,637	*0.86 (0.73-1.01)*
Vitamin E	13	270,107	13,301	*0.95 (0.86-1.06)*
B VITAMINS				
Folic acid	5	131,386	-	*0.89 (0.72-1.09)*
Vitamin B6	3	70,879	7,656	*0.89 (0.83-0.96)*
POLYPHENOLS & FLAVONOIDS				
Green tea	7	99,090	1,577	*0.62 (0.43-0.89)*
Tea	10	314,398	15,389	*0.75 (0.64-0.89)*
FRUITS & VEGETABLES				
Garlic	3	36,604	2,168	*0.69 (0.42-1.13)*
Tomato	11	1,057,923	67,108	*0.74 (0.64-0.85)*
Legumes	8	1,155,076	79,072	*0.89 (0.81-0.97)*
FATTY ACIDS & LIPIDS				
Eicosapentaenoic acid	2	430,507	7,761	*0.95 (0.79-1.15)*
PHYTOESTROGENS				
Soy	8	85,059	5,399	*0.60 (0.41-0.89)*
Genistein	3	70,911	727	*0.78 (0.37-1.63)*
HERBAL & BOTANICAL				
Ginkgo biloba	2	38,308	1,665	*0.93 (0.68-1.27)*
METABOLITES & AMINO ACIDS				
Dietary protein intake	2	109,557	3,064	*0.99 (0.89-1.11)*
DIET				
Plant-based diet	2	211,318	24,516	*0.91 (0.84-0.99)*
DASH Diet	2	152,714	-	*0.93 (0.82-1.06)*
OTHER				
Insoluble dietary fiber intake	3	101,084	-	*0.62 (0.38-1.03)*
Soluble dietary fiber intake	3	80,888	10,851	*0.89 (0.83-0.96)*
Total dietary fiber intake	7	972,077	64,899	*0.92 (0.86-1.00)*
Eggs	3	900,752	53,381	*0.99 (0.96-1.02)*



Exposure group

- Carotenoids
- Vitamins A, C, D, E, K
- B Vitamins
- Polyphenols & Flavonoids
- Fruits & Vegetables
- Fatty Acids & Lipids
- Phytoestrogens
- Herbal & Botanical
- Metabolites & Amino Acids
- Diet
- Other

RR < 1 = negatively associated with prostate cancer risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available)

Include subexposures: No