
# Exposures positively associated with prostate cancer\aggressive risk

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Lutein** | 5 | 3,128 | 1,292 | *1.00 (0.63-1.61)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin D** | 23 | 542,238 | 61,208 | *1.12 (1.06-1.18)* |
| **Vitamin A** | 13 | 70,012 | 17,837 | *1.15 (0.93-1.42)* |
| ***B VITAMINS*** | | | | |
| **Folic acid** | 9 | 91,771 | 19,124 | *1.03 (0.96-1.11)* |
| **Vitamin B12** | 5 | 5,961 | 2,681 | *1.12 (0.97-1.28)* |
| ***ANTIOXIDANTS*** | | | | |
| **Antioxidants** | 3 | 56,601 | 7,271 | *1.15 (0.87-1.50)* |
| **Coenzyme Q10** | 3 | 36,416 | 2,065 | *1.19 (0.79-1.79)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Copper** | 2 | 43,360 | 6,174 | *1.05 (1.00-1.11)* |
| **Iron** | 5 | 184,031 | 11,578 | *1.05 (0.94-1.16)* |
| **Magnesium** | 2 | 43,795 | 6,400 | *1.06 (1.01-1.11)* |
| **Zinc** | 11 | 49,478 | 6,840 | *1.13 (0.85-1.50)* |
| **Calcium** | 24 | 951,593 | 55,176 | *1.14 (1.06-1.23)* |
| ***FERMENTED FOODS & PROBIOTICS*** | | | | |
| **Yogurt** | 3 | 201,324 | 7,680 | *1.25 (1.00-1.56)* |
| **Fermented foods** | 3 | 3,460 | 339 | *1.37 (0.76-2.45)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Docosapentaenoic acid (omega-3)** | 2 | 143,344 | 7,412 | *1.00 (0.97-1.03)* |
| **Omega-3 fatty acids** | 12 | 530,983 | 29,731 | *1.01 (0.96-1.07)* |
| **Polyunsaturated Fat (PUFA)** | 18 | 518,082 | 32,418 | *1.02 (0.95-1.10)* |
| **Alpha-linolenic acid** | 5 | 173,827 | 9,647 | *1.15 (0.92-1.43)* |
| **Monounsaturated Fat (MUFA)** | 7 | 254,582 | 13,412 | *1.22 (1.05-1.42)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Glucosamine** | 3 | 265,129 | 16,136 | *1.06 (1.01-1.11)* |
| **Choline** | 2 | 8,123 | 1,372 | *1.18 (0.83-1.67)* |
| ***DIET*** | | | | |
| **Western pattern diet** | 2 | 44,462 | 1,702 | *1.35 (1.00-1.82)* |
| ***OTHER*** | | | | |
| **Multivitamin** | 4 | 54,340 | 5,633 | *1.01 (0.97-1.04)* |
| **Ultra-processed food intake** | 6 | 248,736 | 13,760 | *1.02 (0.95-1.10)* |
| **Processed meat intake** | 8 | 1,089,615 | 64,702 | *1.09 (1.02-1.17)* |
| **Whole-grain intake** | 3 | 79,458 | 7,209 | *1.11 (0.97-1.26)* |
| **Alcohol** | 25 | 1,826,625 | 45,206 | *1.20 (1.10-1.31)* |
| **Dairy** | 21 | 525,007 | 30,010 | *1.22 (1.10-1.34)* |
| **Red meat** | 10 | 873,126 | 56,254 | *1.60 (1.15-2.23)* |




Exposure group

Carotenoids
Vitamins A, C, D, E, K
B Vitamins
Antioxidants
Minerals & Trace Elements
Fermented Foods & Probiotics
Fatty Acids & Lipids
Metabolites & Amino Acids
Diet
Other

*RR > 1 = positively associated with prostate cancer\aggressive risk. |◆ pooled RR (size reflects number of studies) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | arrowhead = interval continues beyond the axis*

Latest publication per cohort

Include subexposures: Yes