
Exposures negatively associated with bladder cancer risk

Exposure	Estimates	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lutein	3	122,821	1,415	*0.69 (0.38-1.27)*
Beta Carotene	6	368,253	2,305	*0.83 (0.66-1.04)*
Lycopene	3	222,792	1,333	*0.95 (0.90-1.00)*
VITAMINS A, C, D, E, K				
Vitamin A	4	126,541	1,142	*0.72 (0.52-0.98)*
Vitamin C	3	8,606	902	*0.75 (0.56-1.01)*
Vitamin E	5	158,299	1,839	*0.76 (0.58-1.00)*
Vitamin D	9	339,195	4,216	*0.77 (0.34-1.76)*
B VITAMINS				
Folic Acid	5	309,245	2,289	*0.77 (0.61-0.99)*
MINERALS & TRACE ELEMENTS				
Selenium	8	161,440	3,121	*0.73 (0.54-0.97)*
Phosphorus	2	1,147	522	*0.97 (0.27-3.50)*
POLYPHENOLS & FLAVONOIDS				
Tea	9	580,076	9,039	*0.96 (0.79-1.17)*
FRUITS & VEGETABLES				
Legumes	2	2,541	1,268	*0.64 (0.43-0.95)*
Avocado	2	112,328	1,165	*0.78 (0.60-1.01)*
FERMENTED FOODS & PROBIOTICS				
Yogurt	2	101,981	906	*0.64 (0.25-1.66)*
Fermented Foods	5	203,999	2,474	*0.75 (0.50-1.12)*
FATTY ACIDS & LIPIDS				
Omega-3 Fatty Acids	2	1,061	572	*0.31 (0.20-0.48)*
Polyunsaturated Fat PUFA	2	1,061	572	*0.40 (0.24-0.67)*
Omega-6 Fatty Acids	2	1,061	572	*0.46 (0.31-0.69)*
Monounsaturated Fat MUFA	3	649,414	3,842	*0.79 (0.41-1.52)*
OTHER				
Whole Grain Intake	3	219,042	1,666	*0.72 (0.52-1.01)*
Dairy	10	799,763	5,469	*0.83 (0.65-1.05)*
Total Dietary Fiber Intake	2	676,447	3,990	*0.85 (0.76-0.95)*
Alcohol	11	882,294	8,649	*0.86 (0.74-1.00)*



Exposure group

Carotenoids
Vitamins A, C, D, E, K
B Vitamins
Minerals & Trace Elements
Polyphenols & Flavonoids
Fruits & Vegetables
Fermented Foods & Probiotics
Fatty Acids & Lipids
Other

RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis

Latest publication per cohort

Include subexposures: Yes