
# Exposures negatively associated with bladder cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Beta Carotene** | 5 | 366,541 | 1,449 | *0.88 (0.71-1.10)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin C** | 3 | 8,606 | 902 | *0.75 (0.56-1.01)* |
| **Vitamin A** | 2 | 121,519 | 737 | *0.77 (0.39-1.51)* |
| **Vitamin E** | 4 | 157,524 | 1,453 | *0.86 (0.69-1.07)* |
| ***B VITAMINS*** | | | | |
| **Folic Acid** | 4 | 307,597 | 1,465 | *0.73 (0.51-1.05)* |
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Phosphorus** | 2 | 1,147 | 522 | *0.97 (0.27-3.50)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Tea** | 9 | 580,076 | 9,039 | *0.96 (0.79-1.17)* |
| ***FRUITS & VEGETABLES*** | | | | |
| **Legumes** | 2 | 2,541 | 1,268 | *0.64 (0.43-0.95)* |
| **Avocado** | 2 | 112,328 | 1,165 | *0.78 (0.60-1.01)* |
| ***FERMENTED FOODS & PROBIOTICS*** | | | | |
| **Yogurt** | 2 | 101,981 | 906 | *0.64 (0.25-1.66)* |
| **Fermented Foods** | 5 | 203,999 | 2,474 | *0.75 (0.50-1.12)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Monounsaturated Fat MUFA** | 2 | 648,914 | 3,592 | *0.95 (0.44-2.05)* |
| ***OTHER*** | | | | |
| **Whole Grain Intake** | 3 | 219,042 | 1,666 | *0.72 (0.52-1.01)* |
| **Dairy** | 10 | 799,763 | 5,469 | *0.83 (0.65-1.05)* |
| **Total Dietary Fiber Intake** | 2 | 676,447 | 3,990 | *0.85 (0.76-0.95)* |
| **Alcohol** | 10 | 881,889 | 8,536 | *0.86 (0.74-1.00)* |




*RR < 1 = negatively associated with bladder cancer risk. |◆ pooled RR (size reflects number of estimates) |◇ red outline = statistically significant (95% CI excludes 1.0) |◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Latest publication per cohort

Include subexposures: Yes