
Exposures negatively associated with aggressive prostate cancer risk

Exposure	Studies	N	Cases	Pooled RR (95% CI)
CAROTENOIDS				
Lycopene	2	166,781	-	*0.58 (0.39-0.87)*
VITAMINS A, C, D, E, K				
Vitamin E	4	114,280	-	*0.64 (0.40-1.04)*
Vitamin D	7	45,247	-	*0.74 (0.51-1.08)*
MINERALS & TRACE ELEMENTS				
Selenium	5	123,475	-	*0.65 (0.37-1.12)*
Zinc	2	82,482	-	*0.84 (0.17-4.29)*
FRUITS & VEGETABLES				
Tomato	3	124,620	-	*0.70 (0.54-0.90)*
OTHER				
Total dietary fiber intake	2	885,584	5,076	*0.86 (0.63-1.17)*



Exposure group

Carotenoids · Vitamins A, C, D, E, K · Minerals & Trace Elements · Fruits & Vegetables · Other

RR < 1 = negatively associated with aggressive prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line below CI = 95% prediction interval (where available) | arrowhead = interval continues beyond the axis

Include subexposures: Yes

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.