
# Exposures positively associated with aggressive prostate cancer risk

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***MINERALS & TRACE ELEMENTS*** | | | | |
| **Calcium** | 6 | 98,931 | - | *1.53 (0.98-2.38)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Alpha-linolenic acid** | 3 | 383,989 | 3,504 | *1.85 (1.03-3.31)* |
| **Monounsaturated Fat (MUFA)** | 3 | 144,473 | - | *2.39 (1.37-4.15)* |
| **Polyunsaturated Fat (PUFA)** | 6 | 721,149 | - | *2.70 (1.10-6.80)* |
| **Omega-3 fatty acids** | 5 | 720,112 | 4,355 | *3.43 (1.67-7.04)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Betaine** | 3 | 55,116 | 963 | *1.03 (0.60-1.74)* |
| ***DIET*** | | | | |
| **Mediterranean diet** | 2 | 106,146 | - | *1.02 (0.92-1.13)* |
| ***OTHER*** | | | | |
| **Multivitamin** | 2 | 343,481 | 2,040 | *1.27 (1.04-1.56)* |
| **Alcohol** | 4 | 166,503 | - | *1.28 (0.73-2.26)* |
| **Eggs** | 2 | 869,756 | 5,123 | *1.36 (0.87-2.10)* |
| **Dairy** | 3 | 101,926 | 744 | *1.41 (1.14-1.75)* |
| **Processed meat intake** | 4 | 204,990 | 2,302 | *1.41 (1.21-1.65)* |
| **Red meat** | 5 | 129,031 | 1,575 | *1.44 (1.13-1.84)* |




*RR > 1 = positively associated with aggressive prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis*

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.