
Exposures negatively associated with aggressive prostate cancer risk
Meta-analysis of dietary-intake studies

Exposure	Studies	N	Cases	Pooled RR (95% CI)
VITAMINS A, C, D, E, K				
Vitamin E	2	83,022	355	*0.76 (0.29-1.94)*
MINERALS & TRACE ELEMENTS				
Zinc	2	82,482	-	*0.84 (0.17-4.29)*
POLYPHENOLS & FLAVONOIDS				
Tea	2	192,116	-	*0.74 (0.44-1.22)*
FRUITS & VEGETABLES				
Tomato	3	124,620	-	*0.70 (0.54-0.90)*
METABOLITES & AMINO ACIDS				
Betaine	2	54,424	790	*0.81 (0.48-1.38)*
OTHER				
Total dietary fiber intake	2	885,584	5,076	*0.86 (0.63-1.17)*



RR < 1 = negatively associated with aggressive prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval

Include subexposures: Yes

No published subtype baseline: HR ≈ RR ≈ OR approximation. Missing case totals are not estimated.