
# Exposures negatively associated with prostate cancer risk
## Meta-analysis of dietary-intake studies

| Exposure | Studies | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Lutein** | 2 | 1,623 | 677 | *0.65 (0.49-0.86)* |
| **Beta-carotene** | 10 | 94,949 | - | *0.79 (0.63-0.98)* |
| **Lycopene** | 13 | 282,888 | 22,602 | *0.88 (0.79-0.98)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin C** | 10 | 68,446 | 7,637 | *0.86 (0.73-1.01)* |
| **Vitamin E** | 13 | 270,107 | 13,301 | *0.95 (0.86-1.06)* |
| ***B VITAMINS*** | | | | |
| **Folic acid** | 5 | 131,386 | - | *0.89 (0.72-1.09)* |
| **Vitamin B6** | 3 | 70,879 | 7,656 | *0.89 (0.83-0.96)* |
| ***POLYPHENOLS & FLAVONOIDS*** | | | | |
| **Green tea** | 7 | 99,090 | 1,577 | *0.62 (0.43-0.89)* |
| **Tea** | 10 | 314,398 | 15,389 | *0.75 (0.64-0.89)* |
| ***FRUITS & VEGETABLES*** | | | | |
| **Garlic** | 3 | 36,604 | 2,168 | *0.69 (0.42-1.13)* |
| **Tomato** | 11 | 1,057,923 | 67,108 | *0.74 (0.64-0.85)* |
| **Legumes** | 8 | 1,155,076 | 79,072 | *0.89 (0.81-0.97)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Eicosapentaenoic acid** | 2 | 430,507 | 7,761 | *0.95 (0.79-1.15)* |
| ***PHYTOESTROGENS*** | | | | |
| **Soy** | 8 | 85,059 | 5,399 | *0.60 (0.41-0.89)* |
| **Genistein** | 3 | 70,911 | 727 | *0.78 (0.37-1.63)* |
| ***HERBAL & BOTANICAL*** | | | | |
| **Ginkgo biloba** | 2 | 38,308 | 1,665 | *0.93 (0.68-1.27)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Betaine** | 2 | 54,424 | 790 | *0.82 (0.48-1.38)* |
| **Dietary protein intake** | 2 | 109,557 | 3,064 | *0.99 (0.89-1.11)* |
| ***DIET*** | | | | |
| **Plant-based diet** | 2 | 211,318 | 24,516 | *0.91 (0.84-0.99)* |
| **DASH Diet** | 2 | 152,714 | - | *0.93 (0.82-1.06)* |
| ***OTHER*** | | | | |
| **Insoluble dietary fiber intake** | 3 | 101,084 | - | *0.62 (0.38-1.03)* |
| **Soluble dietary fiber intake** | 3 | 80,888 | 10,851 | *0.89 (0.83-0.96)* |
| **Total dietary fiber intake** | 7 | 972,077 | 64,899 | *0.92 (0.86-1.00)* |




Exposure group

Carotenoids
Vitamins A, C, D, E, K
B Vitamins
Polyphenols & Flavonoids
Fruits & Vegetables
Fatty Acids & Lipids
Phytoestrogens
Herbal & Botanical
Metabolites & Amino Acids
Diet
Other

*RR < 1 = negatively associated with prostate cancer risk. | ◆ pooled RR (size reflects number of studies) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | arrowhead = interval continues beyond the axis*

Include subexposures: Yes