
# Exposures positively associated with renal cell carcinoma risk
## Meta-analysis of dietary-intake studies

| Exposure | Estimates | N | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| ***CAROTENOIDS*** | | | | |
| **Lycopene** | 3 | 873,444 | 2,860 | *1.02 (0.73-1.42)* |
| ***VITAMINS A, C, D, E, K*** | | | | |
| **Vitamin A** | 2 | 4,597 | 1,909 | *1.23 (0.53-2.87)* |
| ***B VITAMINS*** | | | | |
| **Folic Acid** | 2 | 4,597 | 1,909 | *1.18 (0.60-2.32)* |
| ***FATTY ACIDS & LIPIDS*** | | | | |
| **Monounsaturated Fat MUFA** | 2 | 777,095 | 1,801 | *1.39 (0.82-2.36)* |
| ***METABOLITES & AMINO ACIDS*** | | | | |
| **Dietary Protein Intake** | 2 | 776,349 | 2,168 | *1.30 (0.75-2.25)* |
| ***OTHER*** | | | | |
| **Processed Meat Intake** | 6 | 1,766,660 | 5,212 | *1.15 (0.94-1.41)* |
| **Red Meat** | 6 | 1,768,446 | 5,800 | *1.43 (1.13-1.81)* |
| **Dairy** | 3 | 3,506 | 1,009 | *2.72 (1.22-6.05)* |




*RR > 1 = positively associated with renal cell carcinoma risk. | ◆ pooled RR (size reflects number of estimates) | ◇ red outline = statistically significant (95% CI excludes 1.0) | ◆ faded = not statistically significant | dashed line = 95% prediction interval | N and case totals sum estimates and may overlap across strata | arrowhead = interval continues beyond the axis*

Include subexposures: Yes